



Europäisches Patentamt  
Eur p an Patent Office  
Office eur p éen d s brevets



Publication number: **0 451 896 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 91200686.3

(22) Date of filing: 25.03.91

(51) Int. Cl.<sup>5</sup>: C12N 1/19, A21D 8/04,  
C12P 19/12, C12N 15/54,  
C12N 15/56, //(C12N1/19,  
C12R1:865)

MOLECULAR BIOLOGY DEPARTMENT  
Do not remove from laboratory

(30) Priority: 28.03.90 EP 90200757  
20.12.90 EP 90203462

(43) Date of publication of application:  
16.10.91 Bulletin 91/42

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: GIST-BROCADES N.V.  
Wateringseweg 1 P.O. Box 1  
NL-2600 MA Delft(NL)

(72) Inventor: Driessen, Marianne  
Vleugelnootlaan 38  
NL-2382 EN Zoeterwoude-Rijndijk(NL)  
Inventor: Osinga, Klaas Anne  
Van der Rijnplantsoen 3  
NL-2253 TC Voorschoten(NL)  
Inventor: Herweijer, Margareta Adriana  
Nieuwe Kelzersgracht 464  
NL-1018 VG Amsterdam(NL)

(74) Representative: Matulewicz, Emil Rudolf  
Antonius, Dr. et al  
Gist-Brocades NV Patents and Trademarks  
Department Wateringseweg 1 P.O. Box 1  
NL-2600 MA Delft(NL)

(64) New yeast strains with enhanced trehalose content, process to obtain such yeasts and the use of these yeasts.

(57) A transformed yeast is disclosed, comprising at least one gene encoding neutral trehalase or trehalose-6-phosphate synthase which gene has been modified such that it differs from a corresponding wild-type gene encoding neutral trehalase or trehalose-6-phosphate synthase, the yeast having as a result a different trehalose content from the untransformed parent yeast. The sugar resistance and drying resistance of the yeast compared to those of the untransformed strain are thereby improved.

EP 0 451 896 A1

hw. trehalose/production/yeast/trehalose-6-phosphate/synthase/  
neutral/trehalase/Saccharomyces/cerevisiae

zie claim 16 ! (ook 19, 20)

## Field of Invention.

This invention relates to yeast, especially baker's yeast.

## 5 Background of the Invention.

It is well known that yeast strains belonging to the genus *Saccharomyces* are capable of fermenting sugars to approximately equal amounts of carbon dioxide and ethanol under anaerobic conditions. The leavening activity of yeast in dough is a result of this activity. The rate of carbon dioxide and ethanol evolution in dough is dependent upon, besides the activity of enzymes involved, the sensitivity of the yeast against the amount of sugar present in the dough. In sugar-rich doughs the leavening activity of the yeast may be much lower than in lean (non-sugared) doughs, caused by a higher osmotic pressure related to the presence of the sugar.

The commercial product baker's yeast is produced in several formulations comprising fresh yeast, dried yeast and frozen yeast. Fresh yeast is available as compressed yeast (27-33% dry matter content) and cream yeast (17-23% dry matter content). Dried yeast is available as active dry yeast (ADY) and as instant dry yeast (IDY) with moisture contents of 6-8% and 3-6%, respectively. Improving the fermentative power of yeast is an ongoing research effort. Both the dried yeast and the moist yeast may be improved to increase their carbon dioxide producing ability in lean (non-sugared) as well as in rich (sugared) doughs so as to reduce leavening time and/or enable the use of less yeast, a considerable cost factor in baking.

Role of trehalose in yeast cells

Trehalose (alpha-D-glucopyranosyl-alpha-D-glucopyranoside) is an important storage compound in vegetative cells and spores of fungi and yeasts. Trehalose appears to serve mainly as a storage carbohydrate during periods of non-proliferation. This applies to the life cycle, the cell cycle and other conditions under which cessation of growth occurs, such as (for example) starvation (Thevelein, J.M. (1984) Microbiol. Rev. 48, 42-59)

Trehalose is accumulated at the end of the reproductive stages and appears to be of importance for the viability of resting cells (Thevelein, *ibid.*). The high trehalose levels in fungal spores are also believed to enhance the resistance of the spores under extreme environmental conditions, such as high and low temperatures and desiccation. Besides a role for trehalose in the viability of spores, several investigators have found a relationship between the level of intracellular or extracellular content of trehalose and stress resistance: the trehalose level is positively correlated with the gassing power of dry yeast (Pollock, G.E., Holstrom, C.D. (1951) Cereal Chem. 28, 498-505). Also a positive correlation of the trehalose level with osmotolerance (Mackenzie, K.F., Sing, K.K., Brown, A.D. (1988), J. Gen. Microbiol. 134, 1661-1666) and with heat tolerance was found (Hottiger, T., Boller, T., Wiemken, A. (1987), FEBS Lett., 220, 113-115). In another study it was found that the level of exogenous, added trehalose correlated with dehydration resistance in stationary phase cells but not in exponential phase cells (Gadd, G.M., Chalmers, K., Reed, R.H. (1987), FEMS Microbiol. Lett., 48, 249-254).

Elevation of the cellular trehalose level.

There is a need to produce yeast of higher quality, such as yeast having higher gassing power. Such a yeast may be obtained by elevating the cellular trehalose content of the yeast. This decreases the harmful effects of several forms of stress on the yeast. Elevation of the cellular trehalose level occurs, under conditions that will lower the growth rate, which can e.g. be achieved by starvation, elevated temperatures and, in general, by forms of stress such as drying and freezing. In yeast manufacturing the lower growth rate can be achieved by lowering the molasses feed at the end of the fermentation. High trehalose levels can also be obtained by applying severe forms of stress as said before. However, these conditions result in yeast with altered properties and lower quality (e.g. viability, gassing power), due to side effects of the applied conditions (Jimenez J., Benitez T., (1988) Curr. Genet. 119, 541-547 and Morris G.J., Winters L., Contson G.E., Clarke K.J. (1983) J. Gen. Microbiol., 129, 2023-2034).

Genetic modification of the trehalose level has been achieved using yeast strains mutated in the RAS2 gene. Yeast RAS proteins are guanine nucleotide-binding proteins and are involved in cAMP formation and possibly also in turnover of inositol phospholipids. Both RAS1 and RAS2 proteins exhibit activities to bind guanine nucleotides as well as to hydrolyze GTP, while this latter activity is abolished in RAS<sup>val19</sup> mutants, in which the nineteenth amino acid (glycine) is changed into valine.

The RAS<sup>val19</sup> mutants exhibit multiple dominant phenotypes which include: (1) failure to accumulate storage carbohydrates such as glycogen and trehalose; (2) inefficient sporulation; (3) sensitivity to nutrient starvation; and (4) sensitivity to heat-shock treatment. Strains which lack the RAS2 gene exhibit virtually opposite phenotypes which include: (1) accumulation of storage carbohydrates and (2) sporulation even on rich media. These mutants fail to grow efficiently on non-fermentable carbon sources and cells homozygous for RAS2 disruptions sporulate on rich media (Tatchell, K., *supra*). These phenotypes are similar to the phenotypes of yeasts affected in the cAMP pathway (Tamanoi, F. (1988) *Biochim. Biophys. Acta.*, **948**, 1-15).

There are conflicting data published concerning the relation between trehalose levels and trehalase activity in the cell.

Specifically, mutations in the RAS2 gene resulted in low levels of neutral trehalase, the trehalose mobilizing enzyme, and elevated levels of accumulated trehalose in early stationary phase cells in the case of a *ras2<sup>-</sup>* mutation (Tatchell, K., Robinson, L.C., Breitenbach, M. (1985), *Proc. Natl. Acad. Sci.*, **82**, 3785-3789). High levels of neutral trehalase and no accumulation of trehalose in stationary phase yeast cells occur with a RAS<sup>2val19</sup> mutation (Toda, T., Uno, I., Ishikawa, T., Powers, S., Kataoka, T., Broek, D., Cameron, S., Broach, J., Matsumoto, K., Wigler, M. (1985), *Cell*, **40**, 27-36). Resistance against different forms of stress was not tested in the mutants mentioned above.

In yet another study, mutants with defects in the cAMP dependent protein kinase cascade were used. A *bcy1* mutation (cAMP independent protein kinase) resulted in a high level of neutral trehalase and low levels of trehalose, while a *cyr1-2(ts)* mutation (thermolabile adenylate cyclase and low levels of cAMP) resulted in a low level of neutral trehalase and a normal level of trehalose, opposite to the results obtained with the *ras2* mutants. (Uno, I., Matsumoto, K., Adachi, K., Ishikawa, T. (1983) *J. Biol. Chem.*, **258**, 10867-10872).

Like the physiologically or environmentally induced modifications of the trehalose content of yeast, such as growth at higher temperatures or starvation conditions, the above mentioned genetically induced modifications have negative side-effects besides elevation of the trehalose content. Therefore modification of the trehalose content genetically via said mutations is not useful for industrial application.

Opposite results about the existence of a direct relationship between the neutral trehalase level and trehalose level in yeast have been presented by different research groups e.g. compare *ras2<sup>-</sup>* mutants, which have a lower neutral trehalase and higher trehalose content, with *cyr1-2(ts)* mutants, which have a lower neutral trehalase but no higher trehalose content.

#### Regulation of the trehalose metabolism in yeast

Trehalose is synthesized from glucose-6-phosphate and UDP-glucose, catalyzed by trehalose-6-phosphate synthase, to form trehalose-6-phosphate, which is processed to trehalose by trehalose-6-phosphate phosphatase. Panek and coworkers (Panek A. C., De Araujo P.S., Moura-Neto V. and Panek A.D. (1987) *Curr. Genet.*, **11**, 459-465) claim that trehalose-6-phosphate synthase activity is inactivated by the cAMP dependent protein kinase. Others, however, claim that trehalose-6-phosphate synthase is not affected by cAMP dependent protein kinase (Vandercammen A., Francois J. and Hers H. (1989) *Eur. J. Biochem.*, **182**, 613-620). Trehalose-6-phosphate synthase activity was measured in stationary phase cells and appeared to be independent of ATP, cAMP and cAMP dependent protein kinase.

Trehalose breakdown is performed by a neutral, cytoplasmic trehalase, which activation is mediated by phosphorylation by the cAMP dependent protein kinase. The non-phosphorylated form is inactive (Uno, I., Matsumoto, K., Adachi, K., Ishikawa, T. (1983), *J. Biol. Chem.*, **258**, 10867-10872). In addition the vacuole contains a so-called acid trehalase which function is not yet known (Harris S.D., Cotter D.A. (1988) *Can. J. Microbiol.* **34**, 835-838).

After a glucose pulse a substantial increase in the intracellular concentration of cyclic-AMP is observed in the first few minutes during the lag phase preceding growth. This is followed by a 6-8 fold increase in the neutral trehalase activity concomitant with the rapid degradation of trehalose. After approximately one hour the neutral trehalase activity is decreased to the low level of activity measured before activation. (Van der Plaats, J.B., Van Solingen, P. (1974) *Biochem. Biophys. Res. Comm.*, **56**, 580-587).

Whereas reduced growth is associated with trehalose accumulation, the induction of growth in resting stages is associated with the rapid mobilization of trehalose due to the activation of trehalase. Addition of a nitrogen source enhances trehalose breakdown and prevents resynthesis (cf. Thevelein (1984) *Microbiol. Rev.*, **48**, 42-59).

The literature data strongly suggest that during periods in which trehalose synthesis is necessary only trehalose-6-phosphate synthase is active while neutral trehalase is inactive and therefore it is prevented that trehalose synthesis and breakdown occur at the same time.

## Summary of the invention.

The present invention provides a transformed yeast comprising at least one gene encoding neutral trehalase or trehalose-6-phosphate synthase which gene has been modified such that it differs from a corresponding wild-type gene encoding neutral trehalase or trehalose-6-phosphate synthase, the yeast having as a result a different trehalose content from the untransformed parent yeast. The transformed yeast of the invention has improved stress resistance compared to the untransformed yeast strain, without the occurrence of the previously mentioned negative side-effects. Furthermore the invention provides an efficient method of trehalose production.

## Detailed description

By 'stress resistance' is meant throughout the present application a higher tolerance to stress conditions e.g. higher osmotic pressure, elevated temperature or conditions of desiccation.

By 'fermentation yield' is meant grams of dry biomass formed per gram of sugars consumed.

By 'neutral trehalase' is meant the cytoplasmic trehalase responsible for the trehalose breakdown.

By 'stringent hybridization' is meant hybridization conditions as described in Maniatis et al. (T. Maniatis, E.E. Fritsch, J. Sambrook (1982), Molecular Cloning, A Laboratory Manual), pp. 387-389.

Surprisingly it has been found now that both neutral trehalase and trehalose-6-phosphate synthase are active when, during the production of baker's yeast, trehalose is synthesized. This is unexpected because during trehalose synthesis, the neutral trehalase is supposed not to be active. This discovery is applied to increase the trehalose content of yeast by modifying the cellular level of trehalase.

According to one aspect of the present invention, the neutral trehalase activity in the yeast cell is lowered by the manipulation of the neutral trehalase gene, thereby increasing the intracellular trehalose content. The thus obtained yeast will have a higher stress resistance.

This approach has the advantage that side effects that have so far occurred simultaneously with trehalose elevation as a result of other genetic or physiological modifications, do not occur in the yeast with a modified cellular neutral trehalase content. Therefore a yeast with an elevated trehalose level due to a modified trehalase content is suitable for industrial use.

According to the invention, yeast strains are provided with improved properties under production and application conditions. With improved properties as used in the specification in connection with the present (modified) yeast strains, is meant higher baking performance, especially improved drying resistance relative to the corresponding untransformed (wild-type) strain and/or higher sugar resistance relative to the corresponding untransformed (wild-type) strain.

Therefore methods are provided for the construction of yeast strains having a higher trehalose content compared to the parent strain. These methods encompass the use of genes which code for enzymes involved in trehalose metabolism, such as trehalose-6-phosphate synthase, trehalose-6-phosphate phosphatase and trehalase. In this way it is possible to regulate the trehalose content of the cell without interference with other metabolic processes (such as described before).

In a preferred embodiment of the invention the gene encoding the neutral trehalase has been isolated. The availability of this gene enables the decrease of the trehalase activity in the cell via the well-known technique of gene-disruption (Rothstein R.J. (1983) in 'Methods in Enzymology' 101, 202) of the respective gene(s). For instance, in a diploid yeast strain one or even both alleles of the trehalase gene can be inactivated (assuming that in a haploid genome, one copy of the trehalase gene is present).

In another embodiment of the invention the neutral trehalase gene can be modified such that a phosphorylation site of the trehalase is modified and phosphorylation cannot occur anymore. It is known from literature that neutral trehalase is activated through phosphorylation, which activation is catalyzed by a cAMP dependent protein kinase. Non-phosphorylated trehalase has a much lower activity. The cyclic AMP-dependent protein kinase consensus recognition site can be modified such that phosphorylation cannot take place anymore. This generates a gene encoding an altered neutral trehalase which cannot be activated anymore via phosphorylation. Any residual activity of the non-phosphorylated trehalase may well be important for the physiology of the cell. This alternative approach therefore may well be the approach of choice in the case that some mobilization of trehalose during biomass formation or in the initial period of baker's yeast performance in dough is essential.

In another embodiment of the invention the gene encoding neutral trehalase is mutagenized at random. Genes coding for trehalase with a lower activity are isolated and used for gene-replacements of the unmodified gene(s) of a yeast strain. Also this approach will lead to an overall lower activity of trehalase and hence a higher level of trehalose.

In another embodiment of the invention the gene encoding neutral trehalase is brought under control of an inducible promoter thereby permitting controlled expression of this gene. This will make it possible to render the gene active under conditions where trehalase activity is beneficial for the cell and to shut off its gene expression under conditions where a stable high trehalose level is wanted, for instance at the end of the yeast biomass production stage.

In another embodiment of the invention the gene coding for neutral trehalase is brought under control of a promoter which is weaker than the promoter of this particular gene.

In still another embodiment of the invention the expression of the neutral trehalase gene(s) can be regulated via the anti-sense RNA approach. In this case, anti-sense RNA of (parts of) the neutral trehalase gene is made in the cell by cloning the coding region (or parts of) in opposite orientation behind a promoter. The RNA synthesized from this gene can base-pair with normal neutral trehalase RNA, thereby preventing its translation. The efficiency with which double stranded RNA is formed ultimately determines the level of neutral trehalase synthesis. Therefore this approach is an alternative to regulate gene expression and is of interest in case the genome contains a family of trehalase genes. Since anti-sense RNA acts *in trans*, such a family of genes can be regulated in a simple way.

In another embodiment of the invention the gene encoding vacuolar acid trehalase is isolated. With this gene as tool this type of trehalase activity can be modified, for instance via gene-disruption techniques, protein engineering, or exchange of promoters (inducible, weaker). This approach is of value when in the absence of any neutral trehalase activity still substantial mobilization of trehalose occurs via the acid trehalase. Therefore, this approach can be used to obtain an optimal level of trehalose in the cell.

In still another embodiment genes encoding enzymes which are directly involved in the synthesis of trehalose, such as trehalose-6-phosphate synthase and trehalose-6-phosphate phosphatase, are cloned. Enhancing the rate of trehalose biosynthesis is another direct approach to increase the level of trehalose. Such an increase can be obtained via genetic engineering techniques applied onto these trehalose biosynthetic genes (increase of gene copy number, protein engineering in order to alter specific activities, use of other promoters and the like).

All these abovementioned embodiments have in common the use of genes coding for enzymes which are directly involved in trehalose metabolism to enhance the trehalose level in the cell. Such an enhanced level is of advantage under conditions of stress, like heat, osmotic pressure, desiccation, frozen dough, and the like.

The yeasts obtained with the above mentioned techniques can be used in strain improvement programs using techniques like DNA mediated transformation techniques as well as techniques like protoplast fusion, mass mating or mutation. Also combinations of these techniques are possible. For example, transformed yeasts can be crossed with another (production) strain. The yeast strains obtained with such improvement programs are included in the scope of the invention.

According to the invention a yeast strain belonging to the genus *Saccharomyces* or to the genus *Kluyveromyces* is used, advantageously a strain belonging to *S. cerevisiae* is applied.

The present invention also provides methods of making doughs (normal and frozen dough), bread and similar products using the yeast of the invention to generate carbon dioxide.

The present invention also provides a yeast with a high content of intracellular trehalose thereby permitting an efficient way of trehalose production.

### Cloning techniques

For general cloning techniques reference is made to the handbook of Maniatis et al. (T. Maniatis, E.F. Fritsch, J. Sambrook (1982) *Molecular Cloning, A Laboratory Manual*). Restriction enzymes are used as recommended by the manufacturer and are obtained either from New England Biolabs (Biolabs), Bethesda Research Laboratories (BRL) or Boehringer Mannheim (Boehringer). In general 1 to 5 units of enzyme are needed to cleave 1 µg of DNA.

Transformation of *E. coli* was carried out using the CaCl<sub>2</sub>-technique (T. Maniatis et al., *supra*).

In further aspects the invention provides a recombinant plasmid or vector which comprises at least part of a neutral trehalase gene and wherein the neutral trehalase gene has been optionally modified in such a way that the said gene codes for trehalase with an activity which is lower than that of the corresponding wild-type gene. It also provides a recombinant plasmid or vector which comprises at least part of a trehalose-6-phosphate synthase gene and wherein the trehalose-6-phosphate synthase gene has been optionally modified in such a way that the said gene codes for trehalose-6-phosphate synthase with an activity which is greater than that of the corresponding wild-type gene.

All publications and patent applications cited in this specification are herein incorporated by reference

as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit and scope of the appended claims.

The following experimental data are given to illustrate the invention. It has to be understood that a person skilled in the art who is familiar with the methods may use other yeast strains and vectors which can be equally used for the purpose of the present invention. These alterations are included in the scope of the invention.

List of deposits (clone, strains and hybridoma cell line)

The following phage clone, strains or cell line have been deposited with CBS (Centraalbureau voor Schimmelcultures Oosterstraat 1, 3742 SA Baarn, the Netherlands, and with ECACC (European Collection of Animal Cell Cultures (ECACC), PHLS-Centre for Applied Microbiology and Re-search, Porton Down, Salisbury, Wiltshire SP4, OJG, UK).

20	lambda gt 11 clone	Accession number	deposition date
25	lambda TRE 16.1.1	CBS 512.90	November 22, 1990
30	Strains	Accession number	deposition date
35	<u>Saccharomyces cerevisiae</u> GRD 11-21 (strain G)	CBS 154.91	March 5, 1991
40	<u>Saccharomyces cerevisiae</u> 227 Ng	CBS 155.91	March 5, 1991
45	<u>Saccharomyces cerevisiae</u> 237 Ng (strain A)	CBS 158.86	March 25, 1986
50	<u>Escherichia coli</u> harbouring plasmid pTRE16.1.1	CBS 115.91	February 19, 1991
55	<u>Escherichia coli</u> harbouring plasmid pTZ19R	CBS 405.87	September 3, 1987
60	Hybridoma cell line producing monoclonal antibody 41-2F10	ECACC 90121901	December 14, 1990

Brief description of the drawing

Figure 1 gives the structure of plasmid pTRE16G418-2. Arrows indicate the direction of transcription.

The orientation of the 16.1.1 segment is not known yet.

Abbreviations: G418-res, Tn5 gene, conferring resistance to G418; ADH promoter, *Saccharomyces cerevisiae* promoter of ADHI gene, linked to G418-res; f1 ori, origin of replication of phage f1; amp<sup>R</sup>, ampicillin resistance gene; 16.1.1, yeast insert as also present in lambda TRE 16.1.1, containing part of the neutral trehalase gene; furthermore, relevant restriction sites have been indicated. The two EcoRI sites at the borders of the 16.1.1 segment are not necessarily present at the corresponding position in the genome of *Saccharomyces cerevisiae*. At least one of these sites results from the way the lambda gt 11 cDNA bank has been constructed. In the innercircle, length is indicated in base pairs.

## 10 EXAMPLE 1

### ACTIVITY OF TREHALASE AND TREHALOSE CONTENTS OF BAKER'S YEAST.

#### Aerobic (fed-batch) fermentation.

15 A yeast cream was prepared using a *Saccharomyces cerevisiae* yeast strain 227 Ng.

A lyophilized culture sample was cultivated in the usual way up to the desired amount of seed yeast. For the cultivation of yeast, laboratory fermentors were used with a content of 10 liters (net volume - 6 liters), provided with a device for a continuous supply of molasses and of an assimilable nitrogen source. 20 The fermentors were also provided with a device for the regulation of the pH and for temperature control. The fermentations were carried out essentially according to G. Reed and H.J. Peppler, Yeast Technology, The AVI Publishing Company Inc., Westport Connecticut, USA, 1973. The cultivation conditions were in particular:

- Molasses mixture, consisting of 80% by weight of beet molasses and 20% by weight of cane molasses, calculated on the basis of 50% sugar;
- Molasses supply (according to Reed and Peppler, page 78/79) according to Table 1;
- The required amount of phosphate was added in the form of mono-ammoniumphosphate, prior to inoculation;
- The required amount of nitrogen was supplied as a 10% solution of NH<sub>3</sub> in water, according to Table 1, below;
- The temperature was kept at 30 °C during the whole fermentation period;
- pH (Reed and Peppler, page 67/68) was kept at 5.0 during the whole fermentation period;
- vitamin B<sub>1</sub> was added in an amount of 6 mg/kg of molasses and 50% sugar at the initial stage of the fermentation.

35 The yeast obtained by these fermentations was concentrated and washed with tap water in a laboratory nozzle centrifuge.

40

45

50

55

Tabl 1

	Tim hours	g. Molasses ad 50% sugar (in total)	g. ammonia ad 10% (in total)
5			
10	0.00	49	20
	1.00	49	20
	2.00	49	20
15	3.00	69	23
	4.00	108	32
	5.00	168	52
20	6.00	246	77
	7.00	325	102
	8.00	404	132
25	9.00	503	166
	10.00	601	200
	11.00	700	228
	12.00	798	228
30	13.00	897	228
	14.00	976	228

35

Enzyme levels were determined in samples taken during the desired fedbatch fermentations.

After filtration, pellets were resuspended in extraction buffer (MES, 25 mM; CaCl<sub>2</sub>, 2.5 mM; PMSF, 1 mM; pH = 7). Cell extracts were made by adding ballotin beads (3 g/ml) and shaking in a vibrax VXR tube shaker 3 times for 3 minutes; cooling between shaking periods (0 °C). Extracts were centrifuged during 10 minutes at 12000 x g and dialyzed overnight against buffer (MES, 5 mM; CaCl<sub>2</sub>, 2.5 mM; pH = 7).

Determination of trehalase activity in the cell free extracts was performed according to Thevelein, J.M., Van Laere, A.J., Beullens, M., Van Assche, J.A., Carlier, A.R. (1983) J. Gen. Microbiol., 129, 716-726, with the exception that tetramethylbenzidine was used as a glucose reagent instead of o-dianisidine. 1 U of trehalase activity is defined as 1 µmol glucose produced per minute.

Trehalose content of the cells was determined with HPLC as follows: fermentation samples were filtered, washed and subsequently boiled. After centrifugation to remove large cell fragments, the supernatant was diluted with acetonitril and water to a final concentration of acetonitril of 50% (v/v). The sample was filtered (0.45 µm) before they were applied to a carbohydrate column with RCSS precolumn and eluted with acetonitril/water, 4:1. Trehalose contents were determined with aid of a refractive index detector and an integrator.

Protein of cell free extracts was determined by the BCA method (Pierce). BSA served as a standard.

The neutral trehalase appeared to be constantly active during the fedbatch fermentation, even when trehalose accumulation starts at approximately 9 hours of fermentation (see Table 2). This result was unexpected because trehalase is supposed to be inactive during trehalose synthesis.

The idea of lowering the cell's neutral trehalase activity was based on this unexpected finding of trehalase being active throughout the fedbatch fermentation.



Table 2

Hours	Trehalose % (w/w)	Trehalase (mU/g dry weight)
2	0.3	5.1
3	0.3	6.2
4	-	6.0
5	0.4	5.4
6	0.1	6.8
7	0.1	6.2
8	0.1	5.9
9	0.3	6.1
10	0.6	7.0
11	1.8	6.3
12	2.7	7.2
13	3.6	7.5
14	4.1	7.8
14.5	5.2	7.7
15	5.1	7.3

Example 2ISOLATION OF PHAGE LAMBDA CLONES CONTAINING PART OF THE GENE CODING FOR NEUTRAL TREHALASE

DNA encoding neutral trehalase was isolated by immuno screening of a lambda gt11 cDNA expression library. Use has been made of a monoclonal antibody (abbreviated 41-2F10) which is directed against the 80 kd subunit of neutral trehalase. This monoclonal has been deposited on December 14, 1990 with the European Collection of Animal Cell Cultures (ECACC), PHLS-Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire SP4, OJG, UK, under the accession number 90121901. This monoclonal antibody reacts with one band of 80 kd on a Western blot prepared from an SDS-PAGE gel onto which a cell free extract of *Saccharomyces cerevisiae* was applied. Also, after incubation of the monoclonal antibody coupled to agarose beads (Afti-Gel Hz hydrazide, Bio-Rad Laboratories, Richmond, Ca, USA) with a cell free extract from *Saccharomyces cerevisiae* neutral trehalase activity is specifically bound to the gel material. After spinning down the beads less than 1% of the neutral trehalase activity is detectable in the supernatant.

The lambda gt11 yeast cDNA library was obtained from Clontech laboratories, Inc. 4030 Fabian way, Palo Alto, California 94903 USA. This library, catalog no. YL 1005b, lot no. 3411, was screened with the monoclonal antibody 41-2F10 according to procedures as described in the Protoblot® immunoscreening system of Promega (catalog no. S3710). See for the screening procedures also Huynh T.V., Young R.A. and Davis R.W. (1985) in: DNA cloning, A practical approach, vol. 1 (D.M. Glover, ed.), pp. 49-78, IRL Press, Oxford.

Several positively reacting clones have been isolated. Lambda TRE16.1.1 has been analyzed further (see below). This phage clone has been deposited with the Centraalbureau voor Schimmelcultures (CBS), Oosterstraat 1, 3742 SA Baarn, the Netherlands on November 22, 1990 under the accession number CBS 512.90.

- 6 An appropriate procedure to propagate the deposited phage clone has been described in Huynh T.V. et al. (see above). E coli strain Y1090 (ATCC no. 37197) has been used as host for propagation of the phage clone (see also Huynh T.V., *ibid.*).

### Example 3

10

#### CONSTRUCTION OF RECOMBINANT PLASMIDS

##### a) pTRE16.1.1

- 15 The yeast insert of lambda TRE16.1.1, has been cloned into the commercially available vector pTZ19R (Pharmacia). This vector has been deposited at the Centraalbureau voor Schimmelcultures, Baarn, The Netherlands, under the accession number 405.87. The cloning procedure was as follows:

- Firstly the yeast insert has been amplified. This was done essentially as described in "PCR Protocols, a guide to methods and applications (Eds. M.A. Innis, D.H. Gelfand, J.J. Sninsky and T.J. White 1990), pp. 253-258", with the following modifications: plaques have been used rather than a phage suspension; the 10 x Taq polymerase buffer contains 20 mM MgCl<sub>2</sub>; the thermal cycling was: 94 °C for 2 minutes, 55 °C for 2 minutes, 72 °C for 3 minutes. The oligonucleotide primers used in this reaction are complementary to DNA sequences surrounding the EcoRI cloning site of the lambda gt11 vector. Their sequences are 5'-GGCGACGACTCCTGGAGCCCG-3' and 5'-CACCAGACCAACTGGTAATGG-3'. After the incubation, the reaction mixture was passed through a BIO-RAD RDP mini column in order to clean the DNA (removal of nucleotides).

- Secondly, the PCR product was digested with EcoRI and cloned into pTZ19R x EcoRI to create plasmid pTRE16.1.1. This clone has been deposited at the CBS under the accession number 115.91 (on February 19, 1991).

30

##### b) Construction of pTRE16G418-2

- In order to provide pTRE16.1.1 with a dominant selection marker, the following ligation has been carried out. A 1.9 kb BglII fragment containing the G418<sup>res</sup> gene under control of the ADHI promoter, was isolated after digestion of plasmid pRBN3 with BglII. pRBN3 has been described fully in European patent application EP-A-306107. This 1.9 kb BglII fragment was cloned into the BamHI site of pTRE16.1.1. This yielded pTRE16G418-2 (see Figure 1).

35

### Example 4

40

#### CONSTRUCTION OF YEAST STRAINS WITH A DISRUPTED NEUTRAL TREHALASE GENE RESULTING IN LOWER NEUTRAL TREHALASE ACTIVITY

- The yeast insert in pTRE16.1.1, when present in an expression vector, directs the synthesis of a polypeptide which is immunologically active against a monoclonal antibody raised against neutral trehalase. The insert has a size of about 1100 bp and originates from a cDNA bank (see Examples 2 and 3). It has been characterized with some restriction enzymes as shown in Figure 1. At this stage it was not known yet from which part of the gene this insert originated. Considering the estimated molecular weight of the neutral trehalase protein on a denaturing gel (about 80 kd (H. App and H. Holzer, J. Biol. Chem. 264, 17583, 1989), the 1100 bp yeast insert most likely does not contain the whole gene.

- According to one embodiment of the invention a yeast strain is provided with a diminished neutral trehalase activity. One way to get such a yeast is to disrupt the gene encoding neutral trehalase. In general, having cloned a gene or part of a gene, there are several methods to disrupt that particular gene in the genome (see Methods in Enzymology, vol. 185, chapter 23: Manipulating yeast genome using plasmid vectors, by T. Stearns, H. Ma and D. Botstein, 1990, Academic Press).

- In one approach, the plasmid-borne copy of the gene (i. . neutral trehalase) is truncated in such a way that it lacks both ends (the true amino terminal and carboxyl terminal regions are missing). Integration of this construct into the chromosome at the neutral trehalase locus leads to two nonfunctional copies of the

gene, one lacking its 5' end, the other its 3' end. This approach has also been followed by us. We reasoned that the 1100 bp insert most likely does not contain the true 5' and 3' end, but only gene-internal sequences. Therefore, we made plasmid pTRE16G418-2, and digested that with NsiI. NsiI cuts twice in plasmid pTRE16G418-2, both times in the part of the neutral trehalase gene (see Figure 1). This digestion generates recombinogenic ends and will direct the integration to the corresponding locus in the genome.

The analysis of the transformants obtained is fully described further below, but at this stage it is already mentioned that the transformants indeed have a diminished neutral trehalase activity. This proves that

1) the yeast insert is part of the neutral trehalase gene and

2) that indeed the yeast insert as present in pTRE16.1.1 (and in pTRE6G418-2) lacks both ends of the neutral trehalase and that it contains a gene-internal segment. In this context we note that we have isolated this yeast insert from a cDNA bank. Almost all the known genes in *Saccharomyces cerevisiae* do not contain introns, and hence we assume that also the neutral trehalase gene is not split. In that case the isolated cDNA clone is colinear with the corresponding genomic gene segment. In case the neutral trehalase gene appears to have introns then still the approach we have followed to disrupt the trehalase gene, will work.

The transformation procedure itself has been carried out as described in European patent application EP-A-306107. Two *Saccharomyces cerevisiae* strains have been used as host: *Saccharomyces cerevisiae* 237 Ng (deposited with the CBS under the accession number 158.86) and *Saccharomyces cerevisiae* GRD 11-21, a diploid strain (deposited with the CBS under the accession number 154.91). For selection of transformants use has been made of the aminoglycoside G418.

These two *Saccharomyces cerevisiae* strains have been transformed with pTRE16G418-2 x NsiI. From each transformation, one transformant has been analysed in further detail. The transformant of strain 237 Ng is abbreviated as ApTRE-dis, the transformant of strain GRD11-21 is abbreviated as GpTRE-dis.

Southern-blot analyses of the genomic DNA of these transformants have confirmed that integration indeed took place at the resident neutral trehalase gene. In both transformants, one allele has been disrupted this way. We note that in the transformants also at least one allele is still unaltered, since both parents are at least diploid (not shown).

Several experiments support the conclusion that indeed the neutral trehalase gene has been disrupted in the transformants (see also Example 5). In one set of experiments, the neutral trehalase activities of parent strain A and its transformant ApTRE-dis has been measured. This was done as follows. Yeasts were grown overnight in shake flasks either in YEP-2.5% glucose or in YEP-2.5% glycerol, 0.2% glucose. Neutral trehalase activity was determined as described in Example 1. The measured activities have been summarized in Table 3. It can be concluded that neutral trehalase activity is significantly lowered in ApTRE-dis, which contains a disrupted neutral trehalase gene. We conclude that the yeast insert, which we have isolated in clone lambda TRE16.1.1 contains part of the gene encoding neutral trehalase.

**Table 3**

Neutral trehalase activity in strain A and its transformant ApTRE-dis (mU/mg cell free extract protein)		
Strain	C-source	
	Glucose	Glycerol + Glucose
A	26.1	27.7
ApTRE-dis	12.2	14.7

The above mentioned example illustrates one possibility to alter the level of neutral trehalase. Alternative methods are possible as well. Some possibilities are described below.

The yeast insert of the isolated phage clone is suitable to perform gene disruption experiments. This

can be carried out as follows: somewhere into such a yeast sequence a DNA segment can be cloned containing a gene that can function as selection marker during in yeast transformation. Examples are the Ura3 gen (in case the recipient host yeast strain has a *ura3*<sup>-</sup> auxotrophy) and so-called dominant selection markers. The latter confer resistance to antibiotics such as gentamycin, hygromycin B and phleomycin or to herbicides such as sulfometuron methyl. Examples are the G418 gene, conferring resistance to gentamycin and brought under control of a *S. cerevisiae* promoter (see for example EP 0306107) and the SMR1-410 gene of *S. cerevisiae*, conferring resistance to sulfometuron methyl (see Casey G., Xiao W. and Rank G.H., J. Inst. Brew. 1988, 94, 93-97). In this way, plasmids can be constructed containing a disrupted trehalase gene, interrupted by selection marker DNA (see above). Such a cassette can then be liberated from the vector backbone and used in gene-transplacement experiments, well described for yeast (Rothstein R.J., supra). In this way, a trehalase gene can be destroyed, resulting in a lower trehalase activity and hence in a higher trehalose content. It will be appreciated to recognize that in the case the host strain contains several trehalase genes, the procedure of gene transplacement can be repeated several times (for instance by using different selection markers). Such altered genes can be integrated into the genome of yeast using the gene-transplacement technique.

With the yeast insert 16.1.1 as probe a genomic library of *Saccharomyces cerevisiae* can be screened. This will ultimately result in the isolation of clones containing the entire neutral trehalase gene and its flanking sequences (promoter/terminator regions). Then, using standard procedures, the natural trehalase promoter can be substituted with a different promoter, preferably a homologous promoter. Such altered genes can be integrated into the genome using gene replacement techniques (Rothstein R.J., supra). This will allow to change the expression of the neutral trehalase gene such that ultimately a lower level of trehalase is being made.

Using standard procedures, the gene can be mutagenized such that it codes for an altered neutral trehalase which cannot be phosphorylated anymore.

It will be understood that the above listed methods are by no way exhaustive.

#### Example 5

#### LEVELS OF TREHALASE ACTIVITY IN STRAINS A AND ApTRE-dis, AS DETERMINED IN CONTINUOUS CULTURE

Since neutral trehalase activity is likely to be regulated by the growth rate of the cells, the effect of disruption of the neutral trehalase gene was determined at two different growth rates. To compare parent and disruption-mutant under identical conditions neutral trehalase activity was measured in cells growing in steady state, in a continuous culture.

Strain A and strain ApTRE-dis were grown in batch for 24 hours in Applikon fermentors (Applikon Dependable Instruments, Schiedam, The Netherlands) with a working volume of 2 litres. Medium and culture conditions were essentially as described by E. Postma, W.A.Scheffers, J.P. van Dijken (1989), Yeast 5, 159-165.

After starting the feed at a dilution rate of 0.06 h<sup>-1</sup> the culture reached steady state in 65 hours, and remained at steady state during 1 day. During that day samples were taken. Then feed was increased to a dilution rate of 0.15 h<sup>-1</sup>. 48 hours later a steady state was reached, and samples were taken.

Activity of neutral trehalase is clearly dependent on the growth rate, as can be seen in Table 4. Both parent and disruption mutant showed higher activities at D = 0.06 h<sup>-1</sup>. At both dilution rates the activity of neutral trehalase of the disruption mutant was about half of the activity of the parent strain. This shows that independent of the growth rate the decrease in activity of neutral trehalase is roughly equivalent with the gene copy number.

Table 4

D (h <sup>-1</sup> )	Trehalase activity (U/g dry weight)	
	A	ApTRE-dis
0.06	11.6	4.8
0.15	7.9	3.3

Example 620 TREHALOSE CONTENTS OF TREHALASE-DISRUPTION MUTANTS OF BAKER'S YEAST

To determine the effect of disruption of a copy of the gene encoding neutral trehalase on the trehalose contents of the cells in the stationary phase the following experiment was performed:

Strains A, G, ApTRE-dis and GpTRE-dis were grown in duplo overnight in 500 ml shake-flasks  
 25 containing 100 ml YEP-medium with 5% glucose. Cells were harvested in the stationary phase, collected and resuspended at t = 0 min. in 70 ml of fresh YEP-medium containing 0.5% glucose. Samples were taken every 90 min. up to 7½ hours.

Trehalose content of the cells was determined with HPLC as follows: fermentation samples were filtered, washed and subsequently boiled. After centrifugation to remove large cell fragments the supernatant was  
 30 injected directly on a Bio-Rad Aminex HPX 87-H column. The column temperature was 65° C, and elution was performed with 0.01 N H<sub>2</sub>SO<sub>4</sub>. Trehalose content was determined with the aid of a refractive index detector and an integrator.

Cell free extracts were prepared as described in Example 1, with the exception that the cell free  
 35 extracts were not dialyzed. Trehalase levels in the cell free extracts were determined as described in Example 1.

Table 5 shows at all time points clear differences between the levels of trehalose in parent and disrupted strains. Strain A initially has a higher trehalose content, which is under the conditions of the experiment less stable than the initially lower trehalose level of strain G.

The activity of neutral trehalase in strain G was clearly lower as compared with the activity in strain A  
 40 (Table 5). Furthermore, in both strain ApTRE-dis and strain GpTRE-dis the trehalase levels were lower with respect to the parent strains (Table 5).

So by lowering the neutral trehalase activity by disruption of the structural gene encoding neutral trehalase it is possible to obtain a yeast with a higher trehalose content. This yeast appeared to be very  
 45 useful for industrial application.

Table 5

Hrs	Trehalose (% w/w)				Trehalase (U/g dry weight)			
	A	ApTRE -dis	G	GpTRE -dis	A	ApTRE -dis	G	GpTRE -dis
0	5.7	7.9	2.6	3.5	4.4	3.0	1.3	0.2
1½	3.7	6.0	2.7	3.6	5.5	3.2	1.3	0.9
3	2.1	5.3	2.7	3.3	3.9	3.8	1.5	1.4
4½	1.2	3.1	2.6	3.5	5.3	4.8	2.1	1.6
6	0.6	2.8	3.1	4.0	5.0	5.0	2.0	1.6
7½	0.1	2.5	3.2	4.3	5.4	4.6	2.7	2.0
Data are mean values of the duplo experiments								

**Claims**

1. A transformed yeast comprising at least one gene encoding neutral trehalase or trehalose-6-phosphate synthase which gene has been modified such that it differs from a corresponding wild-type gene encoding neutral trehalase or trehalose-6-phosphate synthase, the yeast having as a result a different trehalose content from the untransformed parent yeast.
2. A transformed yeast according to claim 1 which has an increased trehalose content compared to the untransformed parent yeast.
3. A transformed yeast according to claim 1 or 2 wherein the said at least one gene encoding neutral trehalase has been modified by disruption of the gene.
4. A transformed yeast according to claim 1 or 2 wherein the expression of the gene encoding neutral trehalase has been modified with the aid of an inducible promoter.
5. A transformed yeast according to claim 1 or 2 wherein the said at least one gene encoding neutral trehalase has been modified such that a phosphorylation site of the trehalase is modified.
6. A transformed yeast according to any one of claims 1 to 5 wherein the at least one gene which has been modified is less active than the corresponding wild-type gene in the breakdown of trehalose, during trehalose synthesis.
7. A yeast obtainable by strain improvement procedures, other than DNA mediated transformation, using a yeast as claimed in any one of claims 1 to 6.
8. A yeast according to any one of claims 1 to 7 wherein the said yeast belongs to the genus Saccharomyces.
9. A cream, compressed, frozen, active dry or instant dry yeast obtainable from a yeast as claimed in any one of claims 1 to 8.

10. A dough or similar product which comprises a yeast as claimed in any one of claims 1 to 9.
11. A process for producing a transformed yeast having a different trehalose content from the untransformed parent yeast, which process comprises modifying at least one gene encoding neutral trehalase and/or trehalose-6-phosphate synthase such that it differs from a wild-type gene encoding neutral trehalase and/or trehalase-6-phosphate synthase.
12. A process according to claim 11 wherein said yeast belongs to the genus Saccharomyces.
13. A yeast according to claim 8 or a process according to claim 12 wherein the genus is Saccharomyces cerevisiae.
14. A process for preparing dough or a similar product, which comprises using transformed yeast as claimed in any one of claims 1 to 9.
15. A process for producing trehalose which comprises using yeast as claimed in any one of claims 1 to 8.
16. A gene coding for neutral trehalase or trehalase-6-phosphate synthase, the gene being in purified and isolated form.
17. A gene or part of a gene coding for neutral trehalase, this part being in purified and isolated form and consisting of
  - i) the DNA insert 16.1.1 in plasmid pTRE16.1.1 (CBS 115.91),
  - ii) a DNA capable of hybridizing under stringent conditions to a DNA of i) or
  - iii) a DNA differing from the DNA's of i) and ii) in codon sequence due to degeneracy of the genetic code.
18. A recombinant plasmid or vector which comprises a neutral trehalase gene and wherein the neutral trehalase gene has been optionally modified in such a way that the said gene codes for trehalase with an activity which is lower than that of the corresponding wild-type gene.
19. A recombinant plasmid or vector which comprises a trehalase-6-phosphate synthase gene and wherein the trehalase-6-phosphate synthase gene has been optionally modified in such a way that the said gene codes for trehalase-6-phosphate synthase with an activity which is greater than that of the corresponding wild-type gene.
20. A microorganism which has been transformed with a vector as claimed in claim 18 or 19.
21. A microorganism according to claim 20 which is a yeast.




Fig. 1





European  
Patent Office

# EUROPEAN SEARCH REPORT

Application Number

EP 91 20 0686

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)		
A	PATENT ABSTRACTS OF JAPAN, vol. 13, no. 549 (C-662)[3897], 7th December 1989; & JP-A-1 225 485 (RES. DEV. CORP. OF JAPAN) 08-09-1989 - - -		C 12 N 1/19 A 21 D 8/04 C 12 P 19/12 C 12 N 15/54 C 12 N 15/56 // (C 12 N 1/19 C 12 R 1:865)		
A	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 262, no. 27, 25th September 1987, pages 13212-13218, The American Society for Biochemistry and Molecular Biology, Inc., US; W. BOOS et al.: "Trehalase of Escherichia coli" * Whole document * - - -				
A,D	JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 264, no. 29, 15th October 1989, pages 17583-17588, The American Society for Biochemistry and Molecular Biology, Inc., US; H. APP et al.: "Purification and characterization of neutral trehalase from the yeast ABYS1 mutant" * Whole document * - - -				
A,D	MICROBIOLOGICAL REVIEWS, vol. 48, no. 1, March 1984, pages 42-59, American Society for Microbiology, Washington, DC, US; J.M. TREVELEIN: "Regulation of trehalose mobilization in fungi" - - -				
A	EP-A-0 306 107 (GIST-BROCADES N.V.) - - - - -				
The present search report has been drawn up for all claims					
Place of search The Hague		Date of completion of search 27 June 91	Examiner VAN PUTTEN A.J.		
<table border="0"><tr><td><b>CATEGORY OF CITED DOCUMENTS</b> X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention</td><td><b>E: earlier patent document, but published on, or after the filing date</b> <b>D: document cited in the application</b> <b>L: document cited for other reasons</b> <hr/><b>&amp;: member of the same patent family, corresponding document</b></td></tr></table>				<b>CATEGORY OF CITED DOCUMENTS</b> X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention	<b>E: earlier patent document, but published on, or after the filing date</b> <b>D: document cited in the application</b> <b>L: document cited for other reasons</b> <hr/> <b>&amp;: member of the same patent family, corresponding document</b>
<b>CATEGORY OF CITED DOCUMENTS</b> X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention	<b>E: earlier patent document, but published on, or after the filing date</b> <b>D: document cited in the application</b> <b>L: document cited for other reasons</b> <hr/> <b>&amp;: member of the same patent family, corresponding document</b>				



## INTERNATIONAL SEARCH REPORT

Inter national Application No  
PCT/EP 94/02167

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6	C12N15/82	C12N15/54	C12N15/11	C12N1/21	A01H5/00
	C12N5/10	C12P19/12	A23L3/3562	A01N3/00	

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H C12P A23L A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	COMPTES RENDUES ACAD. SC. PARIS, vol.259, 20 July 1964 pages 635 - 637 QUILLET, M., ET AL. 'Sur l'accumulation concominante du saccharose et du tréhalose chez plusieurs espèces de Sélaginelles indigènes et exotiques' see the whole document --- -/--	25,35

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

21 November 1994

Date of mailing of the international search report

01.12.94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Maddox, A

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 94/02167

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BIOLOGICAL ABSTRACTS, vol. 90 1990, Philadelphia, PA, US; abstract no. 80013, ADAMS, R.P., ET AL. 'Comparison of free sugars in growing and dessicated plants of Selaginella lepidophylla' see abstract & BIOCHEM. SYST. ECOL., vol.18, no.2/3, 1990 pages 107 - 110 ---	33
X	EUROPEAN JOURNAL OF BIOCHEMISTRY, vol.209, no.3, November 1992 pages 951 - 959 BELL, W., ET AL. 'Characterization of the 56-kDa subunit of yeast trehalose-6-phosphate synthase and cloning of its gene reveal its identity with the product of CIF1, a regulator of carbon catabolite inactivation' see the whole document ---	36
X	YEAST, vol.8, 1992 pages 183 - 192 GONZALES, M.I., ET AL. 'Molecular cloning of CIF1, a yeast gene necessary for growth on glucose' see the whole document ---	36
X	J. BACTERIOLOGY, vol.174, no.3, February 1992 pages 889 - 898 KAASEN, I., ET AL. 'Molecular cloning and physical mapping of the otsBA gene, which encode the osmoregulatory trehalose pathway of Escherichia coli: Evidence that transcription is activated by KatF (AppR)' cited in the application see the whole document ---	36-39
X	EMBL SEQUENCE DATABASE REL. ACC. NO. X69160, 27 May 1993 ---	36-39
X	FEMS MICROBIOL. LETT., vol.107, 1993 page 25-30 MCDOUGALL, J., ET AL. 'A yeast gene for trehalose-6-phosphate synthase and its -complementation of an Escherichia coli otsA mutant' see the whole document ---	36
	--- -/--	

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 94/02167

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 451 896 (GIST-BROCADES) 16 October 1991 see the whole document ---	1-39
A	CURRENT BIOLOGY, vol.2, no.11, 1992 pages 594 - 596 TOMOS, D. 'Life without water' see page 596, left column, last paragraph ---	1-39
A	J. BIOTECHNOLOGY, vol.7, no.1, 1988 pages 23 - 32 COUTINHO, C., ET AL. 'Trehalose as cryoprotectant for preservation of yeast strains' see the whole document ---	32,33
P,X	WO,A,93 17093 (OY ALKO) 2 September 1993 see the whole document ---	36
P,X	EP,A,0 577 915 (ALGIST-BRUGGEMAN) 12 January 1994 see page 7, line 50 - line 54 -----	36

## INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 94/02167

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0451896	16-10-91	AU-B- 636022	08-04-93
		AU-A- 7378291	03-10-91
		JP-A- 5184353	27-07-93
		US-A- 5312909	17-05-94
WO-A-9317093	02-09-93	AU-B- 3500993	13-09-93
		CA-A- 2130082	02-09-93
EP-A-0577915	12-01-94	NONE	